

**EGM 3401**

**Theory Assignment #3**

**Spring 2015**

Due Date: 14 April 2015

## Question 1

Let  $\mathbf{a}$  and  $\mathbf{b}$  be vectors  $\mathbb{E}^3$  and let  $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  is a right-handed orthonormal basis for  $\mathbb{E}^3$ . Furthermore, let  $\mathbf{a} \times \mathbf{b}$  be the vector product between  $\mathbf{a}$  and  $\mathbf{b}$ . Derive an expression for the tensor  $\mathbf{a}^\times \cdot \mathbf{b} = \mathbf{a} \times \mathbf{b}$  in terms of the basis  $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ , that is derive an expression for  $\mathbf{a}^\times$  in the basis  $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  which has the form

$$\mathbf{T} = \sum_{i=1}^3 \sum_{j=1}^3 T_{ij} \mathbf{e}_i \otimes \mathbf{e}_j,$$

where  $\mathbf{T} = \mathbf{a}^\times$ .

## Question 2

Consider a rigid body  $\mathcal{R}$ . Prove the following statements:

- (a)  $m\bar{\mathbf{r}} - \int_{\mathcal{R}} \mathbf{r} dm = \mathbf{0}$ ,
- (b)  $m^{\mathcal{N}}\bar{\mathbf{v}} - \int_{\mathcal{R}} {}^{\mathcal{N}}\mathbf{v} dm = \mathbf{0}$ ,
- (c)  $m^{\mathcal{N}}\bar{\mathbf{a}} - \int_{\mathcal{R}} {}^{\mathcal{N}}\mathbf{a} dm = \mathbf{0}$ ,

where  $(\bar{\mathbf{r}}, {}^{\mathcal{N}}\bar{\mathbf{v}}, {}^{\mathcal{N}}\bar{\mathbf{a}})$ , are, respectively, the position, velocity, and acceleration of the center of mass of  $\mathcal{R}$ .

## Question 3

Let  $\boldsymbol{\tau}$  be a pure torque applied to a rigid body  $\mathcal{R}$ . Prove that the torque  $\boldsymbol{\tau}$  is a free vector and can thus be transported between two points  $P$  and  $Q$  on  $\mathcal{R}$  without changing the torque.

## Question 4

The moment due to a system of forces  $(\mathbf{F}_1, \dots, \mathbf{F}_n)$  and a pure torque  $\boldsymbol{\tau}$  applied to a rigid body  $\mathcal{R}$  relative to a point  $Q$  fixed in  $\mathcal{R}$  is defined as

$$\mathbf{M}_Q = \sum_{i=1}^n (\mathbf{r}_i - \mathbf{r}_Q) \times \mathbf{F}_i + \boldsymbol{\tau}$$

Show that  $\mathbf{M}_Q$  is related to  $\mathbf{M}_P$  (where  $\mathbf{M}_P$  is the moment due the system of forces  $(\mathbf{F}_1, \dots, \mathbf{F}_n)$  and the pure torque  $\boldsymbol{\tau}$  relative to a point  $P$ ) via

$$\mathbf{M}_P = \mathbf{M}_Q + (\mathbf{r}_Q - \mathbf{r}_P) \times \mathbf{F},$$

where  $\mathbf{F}$  is the resultant force acting on the rigid body.

## Question 5

The angular momentum of a rigid body  $\mathcal{R}$  relative to an arbitrary point  $Q$  in an inertial reference frame  $\mathcal{N}$  is defined as

$${}^{\mathcal{N}}\mathbf{H}_Q = \int_{\mathcal{R}} (\mathbf{r} - \mathbf{r}_Q) \times ({}^{\mathcal{N}}\mathbf{v} - {}^{\mathcal{N}}\mathbf{v}_Q) dm$$

Suppose now that the point  $Q$  is equal to a point  $B$  where  $B$  is fixed in  $\mathcal{R}$ . Prove that the angular momentum relative to point  $B$  is given as

$${}^{\mathcal{N}}\mathbf{H}_B = \mathbf{I}_B^{\mathcal{R}} \cdot {}^{\mathcal{N}}\boldsymbol{\omega}^{\mathcal{R}},$$

where  $\mathbf{I}_B^{\mathcal{R}}$  is the moment of inertia tensor of the rigid body  $\mathcal{R}$  relative to point  $B$  and  ${}^{\mathcal{N}}\boldsymbol{\omega}^{\mathcal{R}}$  is the angular velocity of  $\mathcal{R}$  as viewed by an observer in the inertial reference frame  $\mathcal{N}$ .

## Question 6

Let  $\mathcal{R}$  be a rigid body, and let  ${}^{\mathcal{N}}\mathbf{H}_Q$  be the angular momentum of  $\mathcal{R}$  relative to an arbitrary point  $Q$ . Starting with the definition of the angular momentum of a rigid body relative to  $Q$ , that is,

$${}^{\mathcal{N}}\mathbf{H}_Q = \int_{\mathcal{R}} (\mathbf{r} - \mathbf{r}_Q) \times ({}^{\mathcal{N}}\mathbf{v} - {}^{\mathcal{N}}\mathbf{v}_Q) dm,$$

prove that

$${}^{\mathcal{N}}\mathbf{H}_Q = {}^{\mathcal{N}}\tilde{\mathbf{H}} + (\mathbf{r}_Q - \bar{\mathbf{r}}) \times m({}^{\mathcal{N}}\mathbf{v}_Q - {}^{\mathcal{N}}\tilde{\mathbf{v}}).$$

where  ${}^{\mathcal{N}}\tilde{\mathbf{H}}$  is the angular momentum of  $\mathcal{R}$  relative to the center of mass of  $\mathcal{R}$ .

## Question 7

Let  $\mathcal{R}$  be a rigid body and let  $\mathcal{N}$  be an inertial reference frame. Starting with the fundamental form of Euler's second law, that is,

$${}^{\mathcal{N}}\frac{d}{dt} ({}^{\mathcal{N}}\mathbf{H}_O) = \mathbf{M}_O,$$

prove the following two results.

(a) If the reference point is the arbitrary point  $Q$ , then

$${}^{\mathcal{N}}\frac{d}{dt} ({}^{\mathcal{N}}\mathbf{H}_Q) = \mathbf{M}_Q - (\bar{\mathbf{r}} - \mathbf{r}_Q) \times m{}^{\mathcal{N}}\mathbf{a}_Q.$$

(b) If the reference point is the center of mass of  $\mathcal{R}$ , then

$${}^{\mathcal{N}}\frac{d}{dt} ({}^{\mathcal{N}}\tilde{\mathbf{H}}) = \tilde{\mathbf{M}}.$$

The quantities  ${}^{\mathcal{N}}\mathbf{H}_Q$  and  ${}^{\mathcal{N}}\tilde{\mathbf{H}}$  are, respectively, the angular momentum of the rigid body relative to an arbitrary point  $Q$  and the center of mass of  $\mathcal{R}$ .

## Question 8

The kinetic energy of a rigid body is defined as

$$T = \frac{1}{2} \int_{\mathcal{R}} {}^{\mathcal{N}}\mathbf{v} \cdot {}^{\mathcal{N}}\mathbf{v} dm,$$

where the integral is taken over all material points in the body. Prove that  $T$  can be written as

$$T = \frac{1}{2} {}^{\mathcal{N}}\tilde{\mathbf{v}} \cdot {}^{\mathcal{N}}\tilde{\mathbf{v}} + \frac{1}{2} {}^{\mathcal{N}}\tilde{\mathbf{H}} \cdot {}^{\mathcal{N}}\boldsymbol{\omega}^{\mathcal{R}}.$$

This last result is called *Koenig's decomposition* for the kinetic energy of a rigid body.

## Question 9

Consider a system of  $n$  particles with mass  $(m_1, \dots, m_n)$ . Suppose further that the position measured from a point  $O$  fixed in an inertial reference frame  $\mathcal{N}$  can be expressed as  $\mathbf{r}_i = \bar{\mathbf{r}} + \boldsymbol{\rho}_i$ , where  $\bar{\mathbf{r}}$  is the position of the center of mass of the system. In addition, assume that the distance between each of the particles is a constant. Using the definition of angular momentum for the center of mass of a system of particles, show that

$${}^{\mathcal{N}}\tilde{\mathbf{H}} = \left[ \sum_{i=1}^n m_i \{(\boldsymbol{\rho}_i \cdot \boldsymbol{\rho}_i)\mathbf{U} - \boldsymbol{\rho}_i \otimes \boldsymbol{\rho}_i\} \right] \cdot {}^{\mathcal{N}}\boldsymbol{\omega}^{\mathcal{B}},$$

where  $\mathcal{B}$  is the reference frame in which the particles are fixed and  $\mathbf{U}$  is the identity tensor (that is,  $\mathbf{U} \cdot \mathbf{a} = \mathbf{a}$ ).

## Question 10

Let  $\mathcal{R}$  be a rigid body. Show that the angular momentum of the rigid body relative to the center of mass can be written as

$${}^{\mathcal{N}}\tilde{\mathbf{H}} = \left[ \int_{\mathcal{R}} \{(\boldsymbol{\rho} \cdot \boldsymbol{\rho})\mathbf{U} - \boldsymbol{\rho} \otimes \boldsymbol{\rho}\} dm \right] \cdot {}^{\mathcal{N}}\boldsymbol{\omega}^{\mathcal{R}},$$

where  $\mathbf{U}$  is the identity tensor.

## Question 11

Let  ${}^{\mathcal{N}}\mathbf{H}_Q$  be the angular momentum of a rigid body  $\mathcal{R}$  relative to an arbitrary point  $Q$ . Prove that

$${}^{\mathcal{N}}\frac{d}{dt} ({}^{\mathcal{N}}\mathbf{H}_Q) = {}^{\mathcal{R}}\frac{d}{dt} ({}^{\mathcal{N}}\mathbf{H}_Q) + {}^{\mathcal{N}}\boldsymbol{\omega}^{\mathcal{R}} \times {}^{\mathcal{N}}\mathbf{H}_Q.$$